

**Final Exam: Analysis-2 (202200237),** MOD-02-AM: Structures and Systems

Instructors: Carlos Pérez Arancibia and Felix Schwenninger

Date/Time: 29-January-2024, 13:45 – 16:45

- Closed book exam! May use one single-sided handwritten A4-paper.
- All answers must be motivated, including the answers of Section C.
- Answers for Section A could use the four steps (practiced during Tutor Sessions).
- Section Grade:  $\frac{\text{Obtained score}}{\text{Total points}} \times 9 + 1$  (rounded off to one decimal place)
- Course Grade:  $0.4 \times \text{Grade\_Section\_A} + 0.6 \times \text{Grade\_Section\_C}$  (see Assessment Policy for details)
- Good Luck!

**Section C:**

Total Points : 30

1. Evaluate the limit

[10]

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \left| f\left(\frac{k}{n}\right) - f\left(\frac{k-1}{n}\right) \right|$$

where

$$f(t) := \sin(\pi \ln(1+t)).$$

[Hint: Begin by proving the following identity which holds for all functions  $f : [0, 1] \rightarrow \mathbb{R}$  that are continuously differentiable:

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \left| f\left(\frac{k}{n}\right) - f\left(\frac{k-1}{n}\right) \right| = \int_0^1 |f'(t)| dt.$$

Additionally, you could find the inequality  $\sqrt{e} < 2$  helpful.]

2. Consider the function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  given by

$$f(x, y) := \begin{cases} x^2 y \sin\left(\frac{1}{x^2 + y^2}\right) & \text{when } (x, y) \neq (0, 0), \\ 0 & \text{when } (x, y) = (0, 0). \end{cases}$$

(a) Prove from the definition that  $f$  is continuous at  $(0, 0)$ . [3]

(b) Compute the partial derivatives  $\frac{\partial f}{\partial x}(x, y)$  and  $\frac{\partial f}{\partial y}(x, y)$ . [3]

(c) Is  $f$  differentiable at  $(0, 0)$ ? [4]

3. Consider the integral:

$$I(\epsilon) := \iint_{D(\epsilon)} \frac{1}{x+y} dA, \quad 0 < \epsilon < 1,$$

where  $D(\epsilon) \subset \mathbb{R}^2$  is the region bounded by the lines:  $x + y = \epsilon$ ,  $x + y = 1$ ,  $x = 0$ , and  $y = 0$ .

- (a) Evaluate  $I(\epsilon)$  directly using Fubini's theorem. Does the improper integral  $\lim_{\epsilon \rightarrow 0^+} I(\epsilon)$  exist? [3]
- (b) Evaluate  $I(\epsilon)$  using the substitution  $(x, y) = (u - uv, uv)$ . [3]
- (c) Evaluate  $I(\epsilon)$  using the substitution  $(x, y) = (u \cos v, u \sin v)$ . [4]

[Hint: The trigonometric identity  $\sin v + \cos v = \sqrt{2} \sin(v + \frac{\pi}{4})$  could be helpful.]

### Section A:

Total Points : 20

4. (a) Find a closed-form expression for the series [5]

$$\sum_{k=1}^{\infty} \frac{2k}{k+1} (1-x)^k$$

and determine the largest set on which such expression is valid.

- (b) Let  $\alpha \geq 0$  and define  $u_n : [0, \infty) \rightarrow \mathbb{R}$  by [5]

$$u_n(x) = \frac{x^\alpha}{1 + n^2 x^2}, \quad n \in \mathbb{N}.$$

Show that  $\sum_{n=1}^{\infty} u_n(x)$  converges uniformly on  $[0, 1]$  if  $\alpha > 1$ .

5. (a) Prove that if  $f(t)$  is a continuous function on  $[0, 1]$ , then  $\int_0^1 |f(t)| dt = 0$  implies that  $f(t) = 0$  for all  $t \in [0, 1]$ . [5]
- (b) Let  $f(t)$  be a differentiable function on  $[0, 1]$ , and  $f'(t)$  is integrable on  $[0, 1]$ . Assume that  $f(0) = 0$  and  $|f'(t)| \leq |f(t)|$  for all  $t \in (0, 1)$ . Prove that  $f(t) = 0$  for all  $t \in (0, 1)$ . [5]

[Hint: Utilize the Fundamental Theorem of Calculus and consider changing the order of integration over a certain triangular region.]